

Metrics of constant curvature 1 with three conical singularities on 2-sphere

Masaaki Umehara and Kotaro Yamada¹

Osaka Univ.

Kumamoto Univ.

January 28, 1998

Introduction.

Conformal metrics of constant curvature 1 with conical singularities on a closed Riemann surface Σ is bijectively corresponding to branched CMC-1 surfaces in the hyperbolic 3-space with given hyperbolic Gauss map defined on Σ excluded finite points (see [UY2] and also [RUY2]).

It is a fundamental problem to consider the existence of such metrics. Several important results for this direction are found in [T1], [CL], [LT]. Troyanov [T2] gave a classification of metrics of constant curvature 1 with at most two conical singularities on 2-sphere.

In this paper, we shall give a necessary and sufficient condition for the existence and the uniqueness of metric with three conical singularities of given order on 2-sphere. In Section 1, we recall some basic properties of null meromorphic curves in $\mathrm{PSL}(2, \mathbf{C})$. The irreducible case is proved in Section 2 using the study of CMC-1 surfaces developed in [RUY1]. Moreover, applying the same method, we classify all genus zero irreducible CMC-1 surfaces with embedded three regular ends. In Section 3, we give a method for explicit construction of all reducible metric with three singularities.

The authors thank Wayne Rossman and Mikio Furuta for fruitful discussions and encouragement. They are also wishing to thank Yoshinobu Hattori for informative conversations.

1. Preliminaries.

In this section, we recall fundamental properties of null meromorphic curves in $\mathrm{PSL}(2, \mathbf{C})$.

Definition 1.1. Let $F : \Sigma \rightarrow \mathrm{PSL}(2, \mathbf{C})$ be a meromorphic map defined on Riemann surface Σ . Then F is called *null* if

$$\det(F^{-1} \cdot F_z) = 0$$

1991 *Mathematics Subject Classification.* Primary 30F10; Secondary 53C21, 53A10.

¹Supported by Sumitomo Foundation.

holds on Σ , where z is a complex coordinate. (The condition does not depend on the choice of coordinates.)

Let $F : \Sigma \rightarrow \text{PSL}(2, \mathbf{C})$ be a null meromorphic map. We define a matrix α by

$$\alpha = \begin{pmatrix} \alpha_{11} & \alpha_{12} \\ \alpha_{21} & \alpha_{22} \end{pmatrix} := F^{-1} \cdot dF,$$

and set

$$(1.1) \quad g := \alpha_{11}/\alpha_{21}, \quad \omega := \alpha_{21}.$$

Then the pair (g, ω) of a meromorphic function g and a meromorphic 1-form ω on Σ satisfies the equality

$$(1.2) \quad F^{-1} \cdot dF = \begin{pmatrix} g & -g^2 \\ 1 & -g \end{pmatrix} \omega.$$

Conversely, let g be a meromorphic function and ω a meromorphic 1-form on Σ . Then the ordinary differential equation (1.2) is integrable and the solution F is a null map into $\text{PSL}(2, \mathbf{C})$, however F may not be single-valued on Σ . Moreover, F may have essential singularities. We call the pair (g, ω) the *Weierstrass data* of F .

Definition 1.2. Let

$$F = \begin{pmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{pmatrix}$$

be a null meromorphic map of Σ into $\text{PSL}(2, \mathbf{C})$. We call

$$G = \frac{dF_{11}}{dF_{21}} = \frac{dF_{12}}{dF_{22}}$$

the *hyperbolic Gauss map* of F . Furthermore, we call g in (1.1) the *secondary Gauss map* and $Q = \omega dg$ the *Hopf differential* of F .

Let $F : \Sigma \rightarrow \text{PSL}(2, \mathbf{C})$ be a null meromorphic map. Then for $a, b \in \text{SL}(2, \mathbf{C})$, $\hat{F} = a \cdot F \cdot b^{-1}$ is also a null meromorphic map. The associated two Gauss maps $\hat{G}$, $\hat{g}$, and the Hopf differential $\hat{Q}$ of $\hat{F}$ are given by

$$(1.3) \quad \hat{G} = \frac{a_{11}G + a_{12}}{a_{21}G + a_{22}}, \quad \hat{g} = \frac{b_{11}g + b_{12}}{b_{21}g + b_{22}}, \quad \text{and} \quad \hat{Q} = Q,$$

where we set

$$a = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \quad b = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}.$$

Let $(U; z)$ be a complex coordinate of Σ . Now we consider the Schwarzian derivatives $S(G)$ and $S(g)$ of G and g respectively, where

$$(1.4) \quad S(G) = \left[\left(\frac{G''}{G'} \right)' - \frac{1}{2} \left(\frac{G''}{G'} \right)^2 \right] dz^2 \quad \left(' = \frac{d}{dz} \right).$$

The description of the Schwarzian derivative depends on the choice of complex coordinates. However, any difference of two Schwarzian derivatives does not depend on the choice of complex coordinate. The following identity can be checked:

$$(1.5) \quad S(G) = S(g) - 2Q.$$

Conversely, the following fact is known:

Fact 1.3 ([Sm], [UY2]). *Let G and g ($S(G) \neq S(g)$) be non-constant meromorphic functions on a Riemann surface Σ . Then there exists a unique null meromorphic map $F : \Sigma \rightarrow \mathrm{PSL}(2, \mathbf{C})$ with the hyperbolic Gauss map G and the Hopf differential*

$$Q = -\frac{1}{2}(S(G) - S(g)),$$

such that $(g, Q/dg)$ is the Weierstrass data of F .

Moreover, the following fact plays an important role in the latter discussions:

Fact 1.4 ([UY3]). *Let $F : \Sigma \rightarrow \mathrm{PSL}(2, \mathbf{C})$ be a null meromorphic map with hyperbolic Gauss map G and secondary Gauss map g . Then the inverse map F^{-1} is a null meromorphic map with hyperbolic Gauss map g and secondary Gauss map G . In particular, F satisfies the following ordinary differential equation*

$$-F \cdot d(F^{-1}) = dF \cdot F^{-1} = \begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \frac{Q}{dG}.$$

Finally, we point out the following elementary fact on linear algebra:

Fact 1.5 ([RUY1]). *A matrix $a \in \mathrm{SL}(2, \mathbf{C})$ satisfies $a \cdot \bar{a} = \mathrm{id}$ if and only if a is of the form*

$$a = \begin{pmatrix} p & i\gamma_1 \\ i\gamma_2 & \bar{p} \end{pmatrix}, \quad (\gamma_1, \gamma_2 \in \mathbf{R}, \quad p\bar{p} + \gamma_1\gamma_2 = 1).$$

Moreover, a can be diagonalized by a real matrix in $\mathrm{SL}(2, \mathbf{R})$ whenever it is semi-simple.

2. Irreducible metrics with three singularities.

A conformal positive semidefinite metric $d\sigma^2$ on a closed Riemann surface Σ is said to have a *conical singularity* of order β (> -1) at p on Σ if it is asymptotic to the metric of the form $C|z - p|^{2\beta} dz d\bar{z}$ at p , where C is a constant and z is a complex local coordinate of Σ . We denote by $\mathrm{Met}_1(\Sigma)$ the set of (non-vanishing) conformal pseudometrics of constant curvature 1 on Σ with finitely many conical singularities. Let $p_1, \dots, p_n$ be the set of singular points of the given pseudometric $d\sigma^2 \in \mathrm{Met}_1(\Sigma)$ and β_j the order of conical singularity at $z = p_j$. We call a formal sum

$$\beta_1 p_1 + \dots + \beta_n p_n$$

the *divisor* of $d\sigma^2$.

For each metric $d\sigma^2 \in \text{Met}_1(\Sigma)$, there exists a meromorphic function g defined on the universal cover $\tilde{\Sigma}_{p_1, \dots, p_n}$ of $\Sigma_{p_1, \dots, p_n} := \Sigma \setminus \{p_1, \dots, p_n\}$ such that the metric is the pull-back of the canonical metric $d\sigma_0^2 := 4 dz d\bar{z} / (1 + |z|^2)^2$ on $S^2 = \mathbf{C} \cup \{\infty\}$, that is, we have an expression

$$(2.1) \quad d\sigma^2 = g^* d\sigma_0^2 = \frac{4 dg d\bar{g}}{(1 + |g|^2)^2}.$$

Such a function g is unique up to the change

$$(2.2) \quad g \mapsto a \star g, \quad \text{where } a \in \text{PSU}(2) := \text{SU}(2) / \{\pm \text{id}\}.$$

Here, for $a \in \text{PSL}(2, \mathbf{C})$ in general, we set

$$(2.3) \quad a \star g = \frac{a_{11}g + a_{12}}{a_{21}g + a_{22}}, \quad \text{where } a = \pm \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}.$$

We denote by $\pi : \tilde{\Sigma}_{p_1, \dots, p_n} \rightarrow \Sigma_{p_1, \dots, p_n}$ the covering projection. Fix a base point $\tilde{z}_0$ on $\tilde{\Sigma}_{p_1, \dots, p_n}$ and set $z_0 = \pi(\tilde{z}_0)$. For each $z \in \pi^{-1}(z_0)$, there exists a unique deck transformation T such that $T(\tilde{z}_0) = z$. Thus the fundamental group $\pi_1(\Sigma_{p_1, \dots, p_n})$ is identified with the deck transformation group. By (2.1) and (2.2), there exists a representation $\rho_g : \pi_1(\Sigma_{p_1, \dots, p_n}) \rightarrow \text{PSU}(2)$ such that

$$(2.4) \quad g \circ T^{-1} = \rho_g(T) \star g, \quad (T \in \pi_1(\Sigma_{p_1, \dots, p_n})).$$

Later, we will see that the representation ρ_g can be lifted to an $\text{SU}(2)$ -representation.

Metrics $d\sigma^2 \in \text{Met}_1(\Sigma)$ are divided into the three classes defined below;

- (1) A metric is called *irreducible* when the image of the representation ρ_g is not abelian.
- (2) A metric is called $\mathcal{H}^1$ -*reducible* when the image of the representation ρ_g is abelian but non-trivial.
- (3) Then a metric is called $\mathcal{H}^3$ -*reducible* when the image of the representation ρ_g is trivial.

If there exists $a \in \text{PSL}(2, \mathbf{C})$ such that the image of the representation $a \cdot \rho_g \cdot a^{-1}$ is also contained in $\text{PSU}(2)$, then another metric $d\sigma_a^2 := (a \star g)^* d\sigma_0^2$ has the same divisor and the Schwarzian derivative as $d\sigma^2$. Hence, the set

$$(2.5) \quad \{d\sigma_a^2 = (a \star g)^* d\sigma_0^2 \mid a \in \text{PSL}(2, \mathbf{C}) ; a \cdot \text{Im } \rho_g \cdot a^{-1} \subset \text{PSU}(2)\}$$

defines a deformation space of the metric $d\sigma^2 = d\sigma_{\text{id}}^2$ preserving the divisor and the Schwarzian derivative. Since $d\sigma_a^2 = d\sigma^2$ for $a \in \text{PSU}(2)$, the set (2.5) is identified with the set I_Γ in Appendix B, where $\Gamma = \text{Im } \rho_g$.

Suppose that $d\sigma^2$ is $\mathcal{H}^1$ -reducible (resp. $\mathcal{H}^3$ -reducible). Then by Lemma B in Appendix B, the set (2.5) is identified with a totally geodesic subset of dimension one (resp. three) in the hyperbolic 3-space.

We now determine all of irreducible metrics in $\text{Met}_1(S^2)$ with three conical singularities. Reducible case is discussed in the next section.

Theorem 2.1. *There exists an irreducible metric $d\sigma^2$ in $\text{Met}_1(S^2)$ with a given divisor*

$$(2.6) \quad D := \beta_1 p_1 + \beta_2 p_2 + \beta_3 p_3 \quad (\beta_j > -1)$$

if and only if the following inequality holds:

$$(2.7) \quad \cos^2 B_1 + \cos^2 B_2 + \cos^2 B_3 + 2 \cos B_1 \cos B_2 \cos B_3 < 1,$$

where $B_j := \pi(\beta_j + 1)$ ($j = 1, 2, 3$). Moreover, such a metric $d\sigma^2$ is uniquely determined.

Remark 1. The condition (2.7) implies that all β_j ($j = 1, 2, 3$) are not integer. In fact, suppose one of β_j 's, say β_1 , is an integer. Then $\cos B_1 = \pm 1$, and hence (2.7) fails. Conversely, if all β_j ($j = 1, 2, 3$) are not integer, the metric is automatically irreducible (see Corollary 3.2).

Remark 2. Recently, Hattori [H] gave an alternative approach to prove Theorem 2.1 based on the geometry of spherical polytopes.

Remark 3. If a metric $d\sigma^2 \in \text{Met}_1(\Sigma)$ with divisor D in (2.6) is reducible, then Lemma A in Appendix A yields the equality

$$(2.8) \quad \cos^2 B_1 + \cos^2 B_2 + \cos^2 B_3 + 2 \cos B_1 \cos B_2 \cos B_3 = 1.$$

However, even if D satisfies (2.8), it does not imply the existence of the metric. In fact, if $\beta_1, \beta_2, \beta_3 \notin \mathbf{Z}$, such a metric never exists (see Lemma 3.1 and Corollary 3.2).

We identify S^2 with $\mathbf{C} \cup \{\infty\}$ by the stereographic projection, and let z be the canonical complex coordinate of $\mathbf{C}$. For each $d\sigma^2 \in \text{Met}_1(S^2)$, we define the Schwarzian derivative $\tilde{S}(d\sigma^2)$ as

$$(2.9) \quad \tilde{S}(d\sigma^2) := S(g) = \left[\left(\frac{g''}{g'} \right)' - \frac{1}{2} \left(\frac{g''}{g'} \right)^2 \right] dz^2 \quad \left(' = \frac{d}{dz} \right),$$

where g is a meromorphic function defined on $\tilde{S}_{p_1, p_2, p_3}^2$ satisfying (2.1). This is independent of the choice of g . In fact g is unique up to change as in (2.2), and a Möbius transformation (2.3) on g does not change $S(g)$.

Let $d\sigma^2 \in \text{Met}_1(S^2)$ with divisor D as in (2.6), and take a function g as in (2.1). Then for each p_j , there exists $a \in \text{PSU}(2)$ such that

$$(2.10) \quad a \star g = (z - p_j)^{\beta_j+1} (g_0 + g_1 z + \cdots), \quad (g_0 \neq 0).$$

Hence $\tilde{S}(d\sigma^2)$ can be written in the following leading terms in the Laurent expansions at $z = p_j$:

$$(2.11) \quad \tilde{S}(d\sigma^2) = \left[-\frac{\beta_j(\beta_j+2)}{2} \frac{1}{(z-p_j)^2} + \cdots \right] dz^2,$$

and $\tilde{S}(d\sigma^2)$ is holomorphic on S_{p_1, p_2, p_3}^2 . Since the Möbius transformation group acts on the sphere, we may assume

$$(2.12) \quad p_1 = 0, \quad p_2 = 1, \quad \text{and} \quad p_3 = \infty$$

without loss of generality. By (2.11), $\tilde{S}(d\sigma^2)$ is uniquely determined, since the total order of a holomorphic 2-differential on S^2 is 4. In fact, We have

$$(2.13) \quad \tilde{S}(d\sigma^2) = \left[\frac{c_3 z^2 + (c_2 - c_1 - c_3)z + c_1}{z^2(z-1)^2} \right] dz^2,$$

where $c_j = -\beta_j(\beta_j+2)/2 \in \mathbf{R}$ ($j = 1, 2, 3$). Now we set

$$(2.14) \quad G := z, \quad \text{and} \quad Q := \frac{1}{2} \left(\frac{c_3 z^2 + (c_2 - c_1 - c_3)z + c_1}{z^2(z-1)^2} \right) dz^2.$$

By Fact 1.3, there exists a unique null holomorphic map $F_g : \tilde{S}_{p_1, p_2, p_3}^2 \rightarrow \text{PSL}(2, \mathbf{C})$ such that $G \circ \pi$ and g are the hyperbolic Gauss map and the secondary Gauss map respectively. Since $\tilde{S}_{p_1, p_2, p_3}^2$ is simply connected, F_g can be lifted to a null holomorphic map $\tilde{F}_g : \tilde{S}_{p_1, p_2, p_3}^2 \rightarrow \text{SL}(2, \mathbf{C})$. By Fact 1.4, we have the following relations:

$$(2.15) \quad d\tilde{F}_g \cdot \tilde{F}_g^{-1} = \begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \frac{Q}{dG},$$

$$(2.16) \quad g = -\frac{d\tilde{F}_{12}}{d\tilde{F}_{11}} = -\frac{d\tilde{F}_{22}}{d\tilde{F}_{21}},$$

where $\tilde{F}_g = (\tilde{F}_{ij})$. The right-hand side of (2.15) is single-valued and has poles at p_1, p_2 and p_3 . Thus, there exists a representation $\rho_{\tilde{F}_g} : \pi_1(S_{p_1, p_2, p_3}^2) \rightarrow \text{SL}(2, \mathbf{C})$ such that

$$(2.17) \quad \tilde{F}_g \circ T = \tilde{F}_g \cdot \rho_{\tilde{F}_g}(T) \quad (T \in \pi_1(S_{p_1, p_2, p_3}^2)).$$

By (1.3), we have

$$(2.18) \quad \rho_g(T) = \pm \rho_{\tilde{F}_g}(T) \in \text{PSU}(2) \quad (T \in \pi_1(S_{p_1, p_2, p_3}^2)).$$

Here, we consider $\pm a$ ($a \in \text{SU}(2)$) as an element of $\text{PSU}(2) = \text{SU}(2)/\{\pm \text{id}\}$. In particular, $\rho_{\tilde{F}_g}$ is an $\text{SU}(2)$ -representation of $\pi_1(S_{p_1, p_2, p_3}^2)$.

Conversely, if $\tilde{F} = (\tilde{F}_{ij})$ is a solution of (2.15) such that the image of $\rho_{\tilde{F}}$ lies in $\text{SU}(2)$, then (2.18) holds for a meromorphic function g defined by (2.16). So $d\sigma^2$ defined by (2.1) is well-defined on S^2 , that is $d\sigma^2 \in \text{Met}_1(S^2)$, which has divisor D in (2.6). Let T_j be the deck transformation on $\tilde{S}_{p_1, p_2, p_3}^2$ induced from the loop at p_j ($j = 1, 2, 3$). By (2.10) and (2.18), the eigenvalues of $\rho_{\tilde{F}_g}(T_j)$ are $\{\pm e^{iB_j}, \pm e^{-iB_j}\}$, where $B_j = \pi(\beta_j + 1)$. Moreover, we can eliminate this $\pm$ -ambiguity as follows:

Lemma 2.2. *The eigenvalues of $\rho_{\tilde{F}_g}(T_j)$ are $\{-e^{iB_j}, -e^{-iB_j}\}$, where $B_j = \pi(\beta_j + 1)$ ($j = 1, 2, 3$). In particular, $\text{Trace } \rho_F(T_j) = -2 \cos B_j$ holds.*

Proof. Let $\tilde{F}_D$ be the solution of ordinary differential equation

$$\tilde{F}_D \cdot d\tilde{F}_D^{-1} = -d\tilde{F}_D \cdot \tilde{F}_D^{-1} = - \begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \frac{Q}{dG}, \quad \tilde{F}_D(\tilde{z}_0) = \text{id}$$

for the divisor D , where $\tilde{z}_0 \in \tilde{S}_{p_1, p_2, p_3}^2$ is a base point. Then the monodromy representation $\rho_D : \pi_1(S_{p_1, p_2, p_3}^2) \rightarrow \text{SL}(2, \mathbf{C})$ with respect to $\tilde{F}_D$ is conjugate to the representation $\rho_{\tilde{F}_g}$. Let $\tau_j := \text{Trace } \rho_D(T_j) = \pm 2 \cos B_j = \pm 2 \cos \pi(\beta_j + 1)$. Since (2.15) is real analytic in parameters $(\beta_1, \beta_2, \beta_3)$, τ_j 's are real analytic functions in β_j . Here, let $\beta_1 = \beta_2 = \beta_3 = 0$. In this case, $\tilde{F}_D$ is constant because $Q \equiv 0$, and hence each $\rho_D(T_j)$ is the identity matrix. Thus we have

$$\tau_j|_{(\beta_1, \beta_2, \beta_3)=(0,0,0)} = 2 = -2 \cos B_j|_{(\beta_1, \beta_2, \beta_3)=(0,0,0)},$$

and, by real analyticity, $\tau_j = -2 \cos B_j$. Hence $\text{Trace } \rho_{\tilde{F}_g}(T_j) = \text{Trace } \rho_D(T_j) = -2 \cos B_j$, and the eigenvalues of $\rho_D(T_j)$ are $-e^{\pm iB_j}$. (q. e. d.)

Proof of Theorem 2.1. First, we prove the uniqueness of the irreducible metric as follows: Let $d\sigma_l^2 \in \text{Met}_1(S^2)$ ($l = 1, 2$) be two irreducible metrics with the same divisor D as in (2.6). Then we have

$$(2.19) \quad 2Q = \tilde{S}(d\sigma_1^2) = \tilde{S}(d\sigma_2^2).$$

Let g_l be meromorphic functions on $\tilde{S}_{p_1, p_2, p_3}^2$ such that $d\sigma_l^2 = g_l^* d\sigma_0^2$ ($l = 1, 2$). By (2.15), $\tilde{F}_{g_1}$ and $\tilde{F}_{g_2} : \tilde{S}_{p_1, p_2, p_3}^2 \rightarrow \text{SL}(2, \mathbf{C})$ satisfy the same ordinary differential equation. Thus, there exists a matrix $a \in \text{SL}(2, \mathbf{C})$ such that $\tilde{F}_{g_2} = \tilde{F}_{g_1} \cdot a$. So we have

$$(2.20) \quad \rho_{\tilde{F}_{g_2}} = a^{-1} \cdot \rho_{\tilde{F}_{g_1}} \cdot a \quad (a \in \text{SL}(2, \mathbf{C})).$$

Since $\rho_{\tilde{F}_{g_1}}$ and $\rho_{\tilde{F}_{g_2}}$ are both $\text{SU}(2)$ -representations and they are irreducible, we have $a \in \text{SU}(2)$ (see Appendix B). This implies $d\sigma_1^2 = d\sigma_2^2$.

Next, we take $d\sigma^2 \in \text{Met}_1(S^2)$ with divisor D in (2.6). Let T_j ($j = 1, 2, 3$) be a deck transformation corresponding to the loop at $z = p_j$. By Lemma 2.2, the matrix $\rho(T_j)$ ($j = 1, 2, 3$) must have eigenvalues of the form $-e^{\pm iB_j}$. Since $\rho(T_1) \cdot \rho(T_2) \cdot \rho(T_3) = \text{id}$, we have the inequality

$$\cos^2 B_1 + \cos^2 B_2 + \cos^2 B_3 + 2 \cos B_1 \cos B_2 \cos B_3 \leq 1$$




Figure 1: Reflections $\tilde{\mu}_1$, $\tilde{\mu}_2$ and $\tilde{\mu}_3$.

by Lemma A in Appendix A. Since $d\sigma^2$ is irreducible, equality never holds.

Finally, we show the existence of the metric under the condition (2.7).

As (2.12), we identify $S^2_{p_1, p_2, p_3}$ with $\mathbf{C} \setminus \{0, 1\}$. Let μ be the reflection (i.e., conformal transformations reversing orientation) on $\mathbf{C} \setminus \{0, 1\}$ along the real axis. Then there exist reflections $\tilde{\mu}_k$ on the universal cover $\tilde{S}^2_{p_1, p_2, p_3}$ such that

$$\pi \circ \tilde{\mu}_k = \mu \circ \pi, \quad (k = 1, 2, 3),$$

and the deck transformations T_j ($j = 1, 2, 3$) are represented as (see Figure 1)

$$(2.21) \quad T_1 = \tilde{\mu}_1 \circ \tilde{\mu}_2, \quad T_2 = \tilde{\mu}_2 \circ \tilde{\mu}_3, \quad \text{and} \quad T_3 = \tilde{\mu}_3 \circ \tilde{\mu}_1.$$

Let $\tilde{F}$ be a solution of the equation (2.15). Then $\overline{\tilde{F} \circ \tilde{\mu}_k}$ ($k = 1, 2, 3$) is also a solution of (2.15) because of

$$(2.22) \quad \overline{Q \circ \mu} = Q, \quad \text{and} \quad \overline{G \circ \mu} = G.$$

Hence, there exist matrices $\rho_{\tilde{F}}(\tilde{\mu}_k) \in \text{SL}(2, \mathbf{C})$ such that

$$\overline{\tilde{F} \circ \tilde{\mu}_k} = \tilde{F} \cdot \rho_{\tilde{F}}(\tilde{\mu}_k) \quad (k = 1, 2, 3).$$

Since $\tilde{\mu}_k \circ \tilde{\mu}_k = \text{id}$, it holds that

$$(2.23) \quad \rho_{\tilde{F}}(\tilde{\mu}_k) \cdot \overline{\rho_{\tilde{F}}(\tilde{\mu}_k)} = \text{id} \quad (k = 1, 2, 3),$$

and by (2.21), we have

$$(2.24) \quad \begin{aligned} \rho_{\tilde{F}}(\tilde{\mu}_1) \cdot \overline{\rho_{\tilde{F}}(\tilde{\mu}_2)} &= \rho_{\tilde{F}}(T_1), & \rho_{\tilde{F}}(\tilde{\mu}_2) \cdot \overline{\rho_{\tilde{F}}(\tilde{\mu}_3)} &= \rho_{\tilde{F}}(T_2), \\ \text{and} \quad \rho_{\tilde{F}}(\tilde{\mu}_3) \cdot \overline{\rho_{\tilde{F}}(\tilde{\mu}_1)} &= \rho_{\tilde{F}}(T_3). \end{aligned}$$

If there exists a solution $\tilde{F}$ of (2.15) such that

$$(2.25) \quad \rho_{\tilde{F}}(\tilde{\mu}_k) \in \text{SU}(2) \quad (k = 1, 2, 3),$$

it follows that $\rho_{\tilde{F}}(T_j) \in \text{SU}(2)$ for $j = 1, 2, 3$ because of (2.24). For such an $\tilde{F}$, we set g as in (2.16) and $d\sigma^2$ as in (2.1). Then $\rho_g(T) \in \text{SU}(2)$ for each deck transformation T on $\tilde{S}_{p_1, p_2, p_3}$. This implies that $d\sigma^2 \in \text{Met}_1(S^2)$. Moreover, the divisor of $d\sigma^2$ is D by (1.5).

Thus, it is enough to show that there exists a solution $\tilde{F}$ of (2.15) which satisfy (2.25). To do this, we use the following argument similar to the proof of Proposition 6.7 and Proposition 5.6 in [RUY1].

Step 1: Let z_0 be a point on the segment $(-\infty, 0)$ of the real axis on $\mathbf{C}$ and take a point $\tilde{z}_0 \in \tilde{S}_{p_1, p_2, p_3}$ such that $\pi(\tilde{z}_0) = z_0$ and $\tilde{\mu}_1(\tilde{z}_0) = \tilde{z}_0$. Take the solution $\tilde{F}$ of (2.15) with an initial condition $\tilde{F}(\tilde{z}_0) = \text{id}$. By (2.22), $\tilde{F} \circ \tilde{\mu}_1$ is also a solution of (2.15). Moreover, since $\tilde{\mu}_1(\tilde{z}_0) = \tilde{z}_0$, $\tilde{F} \circ \tilde{\mu}_1$ has the same initial condition as $\tilde{F}$. Hence we have $\tilde{F} \circ \tilde{\mu}_1 = \tilde{F}$, and then $\rho_{\tilde{F}}(\tilde{\mu}_1) = \text{id}$.

Step 2: Let $\tilde{F}$ be as in the previous step. By (2.24) and Lemma 2.2, the eigenvalues of $\rho_{\tilde{F}}(\tilde{\mu}_2)$ are $-e^{\pm iB_2}$. Moreover, $\sin B_2 \neq 0$ because β_2 is not an integer. In particular, $\rho_{\tilde{F}}(\tilde{\mu}_2)$ is semisimple. By Fact 1.5 and (2.23), there exists a matrix $u \in \text{SL}(2, \mathbf{R})$ such that

$$u^{-1} \cdot \rho_{\tilde{F}}(\tilde{\mu}_2) \cdot u = \begin{pmatrix} -e^{iB_2} & 0 \\ 0 & -e^{-iB_2} \end{pmatrix}.$$

Let $\hat{F} := \tilde{F} \cdot u$. Then $\hat{F}$ is also a solution of (2.15) and

$$(2.26) \quad \begin{aligned} \rho_{\hat{F}}(\tilde{\mu}_1) &= u^{-1} \cdot \rho_{\tilde{F}}(\tilde{\mu}_1) \cdot u = u^{-1} \cdot \rho_{\tilde{F}}(\tilde{\mu}_1) \cdot u = \text{id}, \\ \rho_{\hat{F}}(\tilde{\mu}_2) &= u^{-1} \cdot \rho_{\tilde{F}}(\tilde{\mu}_2) \cdot u = u^{-1} \cdot \rho_{\tilde{F}}(\tilde{\mu}_2) \cdot u = \begin{pmatrix} -e^{iB_2} & 0 \\ 0 & -e^{-iB_2} \end{pmatrix} \end{aligned}$$

hold because u is a real matrix.

Step 3: Let $\hat{F}$ be as in Step 2. By Fact 1.5 and (2.23), $\rho_{\hat{F}}(\tilde{\mu}_3)$ can be written as

$$\rho_{\hat{F}}(\tilde{\mu}_3) = \begin{pmatrix} q & i\delta_1 \\ i\delta_2 & \bar{q} \end{pmatrix}, \quad (\delta_1, \delta_2 \in \mathbf{R}, \quad q\bar{q} + \delta_1\delta_2 = 1).$$

Then by (2.24) and Lemma 2.2, we have

$$q = \frac{i}{\sin B_1} (\cos B_2 + e^{iB_1} \cos B_3).$$

Hence, by the assumption (2.7),

$$\begin{aligned} \delta_1\delta_2 &= 1 - q\bar{q} \\ &= 1 - \frac{\cos^2 B_1 + \cos^2 B_2 + \cos^2 B_3 + 2 \cos B_1 \cos B_2 \cos B_3}{\sin^2 B_1} > 0. \end{aligned}$$

Let

$$\check{F} = \hat{F} \cdot \begin{pmatrix} (\delta_1/\delta_2)^{1/4} & 0 \\ 0 & (\delta_2/\delta_1)^{1/4} \end{pmatrix} \in \text{SL}(2, \mathbf{R}).$$

Then $\check{F}$ is a solution of (2.15) and

$$\rho_{\check{F}}(\tilde{\mu}_1) = \text{id}, \quad \rho_{\check{F}}(\tilde{\mu}_2) = \begin{pmatrix} -e^{iB_1} & 0 \\ 0 & -e^{-iB_1} \end{pmatrix}, \quad \rho_{\check{F}}(\tilde{\mu}_3) = \begin{pmatrix} q & i\delta \\ i\delta & \bar{q} \end{pmatrix} \in \text{SU}(2),$$

where $\delta = (\delta_1 \delta_2)^{1/2}$. Thus, we have a desired metric $d\sigma^2$ induced from $g = -d\check{F}_{12}/d\check{F}_{11}$. (q. e. d.)

Metrics with conical singularities are closely related to CMC-1 (constant mean curvature 1) surfaces in the hyperbolic 3-space H^3 . In fact, as shown in [UY2, Theorem 2.2] the set $\text{Met}_1(\Sigma)$ corresponds bijectively to the set of branched CMC-1 immersions of Σ excluded finite points of finite total curvature with prescribed hyperbolic Gauss map. The one direction of the correspondence is given as follows: Let $x : M := \Sigma \setminus \{p_1, \dots, p_n\} \rightarrow H^3$ be a conformal CMC-1 immersion whose induced metric ds^2 is complete and of finite total curvature. We set

$$d\sigma_x^2 := (-K)ds^2,$$

where K is the Gaussian curvature of ds^2 . Then it can be extended to a pseudo-metric on Σ and $d\sigma_x^2 \in \text{Met}_1(\Sigma)$ holds (cf. [B]). The converse correspondence is described in [UY2, Section 2].

Definition 2.3. A regular end of a CMC-1 immersion x is called Type I, if the Hopf differential of the immersion has pole of order 2 at the end (cf. [RUY2]).

As seen in [UY1, Section 5], *a regular end of the CMC-1 immersion x is asymptotic to a certain catenoid cousin end if and only if the end is of Type I and embedded.*

Using the same argument of the proof of Theorem 2.1, we can classify the set of irreducible CMC-1 surface in the hyperbolic 3-space of genus zero, with three ends asymptotic the catenoid cousins.

Theorem 2.4. *If a triple of real numbers $(\beta_1, \beta_2, \beta_3)$ satisfy (2.7) for $B_j = \pi(\beta_j + 1)$ ($j = 1, 2, 3$), there exists a unique irreducible conformal CMC-1 immersion $x : S_{p_1, p_2, p_3}^2 \rightarrow H^3$ such that all ends p_1, p_2, p_3 are of embedded and Type I, and the order of the pseudometric $d\sigma_x^2$ at p_j is β_j . Conversely, any conformal irreducible immersed CMC-1 surface of genus zero with three embedded Type I ends are obtained in such a manner.*

Proof. In fact, such a surface is realized by an conformal CMC-1 immersion $x : \mathbf{C} \setminus \{0, 1\} \rightarrow H^3$ with the following properties:

- (1) Since three ends are of Type I, the Hopf differential Q of x has poles of order 2 at the ends 0, 1 and ∞ . Then necessarily it has two zeros q_1, q_2 of order 1 on $\mathbf{C} \setminus \{0, 1\}$. (When Q has the only zero of order two, we regard $q_1 = q_2$.)
- (2) Since all ends are regular and embedded, the hyperbolic Gauss map G of f has two branch point of order 1 at zeroes of Q , and no branch point elsewhere (see [UY3]). (Hence, G is a meromorphic function of degree 2.)

(3) $\tilde{S}(d\sigma_x^2) - S(G) = 2Q$ holds (see [UY2, (2.3)]).

(4) Since the order of the pseudometric $d\sigma_x^2$ at the umbilic points q_1, q_2 are equal to the order of zeroes of Q (see [UY3]), $d\sigma_x^2$ has the divisor of the form

$$D' := \beta_1 p_1 + \beta_2 p_2 + \beta_3 p_3 + q_1 + q_2, \quad (\beta_j > -1).$$

The metric $d\sigma_x^2$ is irreducible if and only if (2.7) holds (cf. Appendix A).

We may assume that, $p_1 = 0$, $p_2 = 1$ and $p_3 = \infty$. By (1), (2) and (3), the top term of $2Q$ of Laurent expansion at $z = p_j$ is the same as that of $\tilde{S}(d\sigma_x^2)$. Thus we have

$$Q = \frac{1}{2} \left(-\frac{\beta_j(\beta_j + 2)}{2} \frac{1}{(z - p_j)^2} + \dots \right) dz^2 \quad (j = 1, 2, 3).$$

Since the Hopf differential Q is holomorphic on S_{p_1, p_2, p_3}^2 , we have

$$(2.27) \quad Q = \frac{1}{2} \left(\frac{c_3 z^2 + (c_2 - c_1 - c_3)z + c_1}{z^2(z - 1)^2} \right) dz^2,$$

where $c_j = -\beta_j(\beta_j + 2)/2 \in \mathbf{R}$ ($j = 1, 2, 3$). By (2), $G^*d\sigma_0^2$ has the divisor of the form

$$D_G := q_1 + q_2.$$

Since the hyperbolic Gauss map G has an ambiguity of Möbius transformations, we can set

$$(2.28) \quad G = z + \frac{(q_1 - q_2)^2}{2\{2z - (q_1 + q_2)\}}.$$

Then by [RUY1, Theorem 3.1], we can see the uniqueness of irreducible CMC-1 immersion x with the hyperbolic Gauss map G and the Hopf differential Q . So it is sufficient to show the existence of such a surface. The following proof is almost same as that of Theorem 2.1: Let μ be the reflection with respect to the real axis and take the reflections $\tilde{\mu}_j$ ($j = 1, 2, 3$) on the universal cover $\tilde{S}_{p_1, p_2, p_3}^2$ as in the proof of Theorem 2.1 (see Figure 1).

Let $\tilde{F}$ be a solution of the equation

$$(2.29) \quad d\tilde{F} \cdot \tilde{F}^{-1} = \begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \frac{Q}{dG}$$

for G in (2.28) and Q in (2.27). Then $\overline{\tilde{F} \circ \tilde{\mu}_k}$ ($k = 1, 2, 3$) is also a solution of (2.15) because of

$$(2.30) \quad \overline{Q \circ \mu} = Q, \quad \text{and} \quad \overline{G \circ \mu} = G.$$

Hence, there exist matrices $\rho_{\tilde{F}}(\tilde{\mu}_j) \in \text{SL}(2, \mathbf{C})$ such that

$$(2.31) \quad \overline{\tilde{F} \circ \tilde{\mu}_j} = \tilde{F} \cdot \rho_{\tilde{F}}(\tilde{\mu}_j) \quad (j = 1, 2, 3).$$

Now by the completely same argument as in the proof of Theorem 2.1, there exists a solution $\tilde{F}$ of (2.29) such that $\rho_{\tilde{F}}(T_j) \in \text{SU}(2)$ for $j = 1, 2, 3$. For such an $\tilde{F}$, we

set g as in (2.16) and $d\sigma^2$ as in (2.1). Then $d\sigma^2 \in \text{Met}_1(S^2)$ has the divisor D' . By [UY2, Theorem 2.2], there exists a branched CMC-1 immersion $x : S^2_{p_1, p_2, p_3} \rightarrow H^3$ whose hyperbolic Gauss map and Hopf differential are G and Q respectively such that $d\sigma_x^2 = d\sigma^2$. One can easily check that the metric given by

$$ds^{2\sharp} := (1 + |G|^2)^2 \left| \frac{Q}{dG} \right|^2$$

is positive definite and complete. Thus by [RUY1, Lemma 2.3], so is the first fundamental form ds^2 . Hence x is the desired one. (q. e. d.)

Proposition 2.5. *Let $x : S^2_{p_1, p_2, p_3} \rightarrow H^3$ be complete CMC-1 surface with three ends of Type I. Then the total absolute curvature TA of x is greater than or equal to 4π .*

Remark. In [UY1], the authors showed $\text{TA} > 2\pi$ for three ended CMC-1 surfaces. The estimate in Proposition 2.5 is sharper than this.

Proof. The associated pseudometric $d\sigma_x^2$ has the divisor $D' := \beta_1 p_1 + \beta_2 p_2 + \beta_3 p_3 + q_1 + q_2$ where q_1, q_2 are umbilic points of x . Then we have

$$\begin{aligned} \frac{1}{2\pi} \text{TA} &= \frac{1}{2\pi} \int_{S^2_{p_1, p_2, p_3}} (-K) dA_{ds^2} = \frac{1}{2\pi} \int_{S^2_{p_1, p_2, p_3}} dA_{d\sigma^2} \\ &= \chi(S^2) + |D| = 4 + \beta_1 + \beta_2 + \beta_3. \end{aligned}$$

On the other hand, the metric $d\sigma_x^2$ induces a monodromy representation $\rho_g : \pi_1(S^2_{p_1, p_2, p_3}) \rightarrow \text{PSU}(2)$. As seen in the proof of Theorem 2.1, it can be lifted to a representation $\rho_{\tilde{F}_g} : \pi_1(S^2_{p_1, p_2, p_3}) \rightarrow \text{SU}(2)$. Let T_j ($j = 1, 2, 3$) be the deck transformation corresponding the the loop surrounding p_j . Then the eigenvalues of $\rho_{\tilde{F}_g}$ is $-e^{\pm i B_j}$ where $B_j := \pi(\beta_j + 1)$, which can be proved by the same method as in Lemma 2.2. By Lemma A in Appendix A, we have

$$\cos^2 B_1 + \cos^2 B_2 + \cos^2 B_3 + 2 \cos B_1 \cos B_2 \cos B_3 < 1.$$

Then it can be easily seen that $B_1 + B_2 + B_3 \geq \pi$, which yields $\beta_1 + \beta_2 + \beta_3 \geq -2$. (q. e. d.)

3. Reducible metrics with three singularities.

In this section, we give a necessary and sufficient condition for the existence of reducible metrics with given divisors. As in the previous section, we identify $S^2 = \mathbf{C} \cup \{\infty\}$, and set $(p_1, p_2, p_3) = (0, 1, \infty)$.

Lemma 3.1. *Let $d\sigma^2 \in \text{Met}_1(S^2)$ be an $\mathcal{H}^3$ -reducible (resp. $\mathcal{H}^1$ -reducible) pseudometric with divisor D as in (2.6). Then all of β_j 's are integers (resp. exactly one of β_j 's is an integer). Namely, at least one of β_j is an integer for reducible metrics.*

Proof. Let $d\sigma^2 \in \text{Met}_1(S^2)$ be an $\mathcal{H}^3$ -reducible metric with divisor D as in (2.6). Then, by definition, the representation ρ_g as in (2.4) is trivial. Hence g satisfying (2.1) is a single-valued meromorphic function on $S^2_{p_1, p_2, p_3}$. Since p_j ($j = 1, 2, 3$) are conical singularities of $d\sigma^2$, g can be extended to a meromorphic function on S^2 . Hence β_j ($j = 1, 2, 3$) are integers.

Next, assume $d\sigma^2$ is $\mathcal{H}^1$ -reducible and all β_j 's are non-integral numbers. Since $d\sigma^2$ is reducible, we can choose g as in (2.1) such that $\rho_g(T)$ are diagonal for all $T \in \pi_1(S^2_{p_1, p_2, p_3})$. Then $g \circ T_k = e^{2\pi i(\beta_k + 1)}g$ because of (2.10). Hence $g_1 := z^{-\beta_1 - 1}g$ is single-valued on $\tilde{S}^2_{p_2, p_3}$, and $g_2 := (z - 1)^{-\beta_2 - 1}g_1$ is single-valued on $\tilde{S}^2_{p_3}$. Since $S^2_{p_3}$ is simply-connected, g_2 is single-valued on S^2 and g can be written as

$$(3.1) \quad g = z^\mu (z - 1)^\nu \frac{a(z)}{b(z)} \quad (\mu, \nu \in \mathbf{R} \setminus \mathbf{Z}),$$

where $a(z)$ and $b(z)$ are mutually prime polynomials whose roots are distinct from 0 and 1. Then we have

$$(3.2) \quad \begin{aligned} dg &= z^{\mu-1} (z - 1)^{\nu-1} \frac{p(z)}{q(z)} dz, \\ p(z) &:= \{\nu z + \mu(z - 1)\}a(z)b(z) + z(z - 1)\{a'(z)b(z) - a(z)b'(z)\}, \\ q(z) &:= \{b(z)\}^2. \end{aligned}$$

Since $p(0) = -\mu a(0)b(0)$, $p(1) = \nu a(0)b(0)$, $q(0) = \{b(0)\}^2$ and $q(1) = \{b(1)\}^2$ are not equal to 0, the roots of p and q are distinct from 0 and 1. Moreover, p and q are mutually prime. In fact, assume there exists a common root ξ of p and q . Then $b(\xi) = 0$, and by assumption, $\xi \neq 0, 1$ and $a(\xi) \neq 0$. Then $0 = p(\xi) = \xi(\xi - 1)a(\xi)b'(\xi)$ implies $b'(\xi) = 0$. Hence ξ is a multiple root of b . Let $b(z) = (z - \xi)^m \tilde{b}(z)$, where $m \geq 2$ be an integer and $\tilde{b}(z)$ is a polynomial such that $\tilde{b}(\xi) \neq 0$. Then we have

$$dg = z^{\mu-1} (z - 1)^{\nu-1} \frac{(z - \xi)r(z) - z(z - 1)\tilde{b}(z)}{(z - \xi)^{m+1}\tilde{b}(z)^2} dz,$$

where $r(z) = (\nu z + \mu(z + 1))a\tilde{b} + z(z - 1)(a'\tilde{b} - a\tilde{b}')$ is a polynomial in z . Since $m \geq 2$, ξ is a ramification point of g , and then, $d\sigma^2$ has a conical singularity at ξ , which is a contradiction.

Thus p and q are mutually prime whose roots are distinct from 0 and 1. If p has a root η , then η is a ramification point of g , and then, a conical singularity of $d\sigma^2$. Hence $p(z)$ must be a constant. By (3.2), p is formally a polynomial of degree $\deg a + \deg b + 1$. Then the highest term must vanish: $\mu + \nu + \deg a - \deg b = 0$. This shows that the order of dg at $z = \infty$ must be integer, and then β_3 is an integer. This is contradiction, and hence at least one of β_j 's must be integer.

On the other hand, let $d\sigma^2 \in \text{Met}_1(S^2)$ be $\mathcal{H}^1$ -reducible and exactly one of β_j 's is a non-integral number. Without loss of generality, we assume β_1 is non-integral. Take g as in (2.1). Since β_2 and β_3 are integers, g is well-defined on the universal

cover of $S_{p_1}^2$. Here $S_{p_1}^2$ is simply connected. Then g is single-valued on S^2 itself, and hence, g is a meromorphic function on S^2 . This shows that β_1 is an integer, contradiction.

Hence, if $d\sigma^2 \in \text{Met}_1(S^2)$ with divisor D in (2.6) is $\mathcal{H}^1$ -reducible, exactly one of β_j 's is an integer. (q. e. d.)

As an immediate consequence, we have the following corollary.

Corollary 3.2. *Let $d\sigma^2 \in \text{Met}_1(S^2)$ has exactly three singularities with orders β_1, β_2 and β_3 . Then the following three assertions are true.*

- (1) $d\sigma^2$ is $\mathcal{H}^3$ -reducible if and only if all of β_j 's are integers.
- (2) $d\sigma^2$ is $\mathcal{H}^1$ -reducible if and only if exactly one of β_j 's is an integer.
- (3) $d\sigma^2$ is irreducible if and only if all of β_j 's are non-integers.

Remark. For a metric in $\text{Met}_1(S^2)$ with more than three singularities, such a simple criterion for reducibility is not expected: There exists a reducible metric $d\sigma^2 \in \text{Met}_1(S^2)$ with divisor

$$D' = \beta_1 p_1 + \beta_2 p_2 + \beta_3 p_3 + q_1 + q_2$$

such that all β_j 's are non-integral numbers. In fact,

$$g = c z^\mu (z-1)^\nu (z-a) \quad (c \in \mathbf{C} \setminus \{0\}, a \in \mathbf{C} \setminus \{0, 1\})$$

induces such a metric whenever $\mu + \nu$ is not an integer. On the other hand, we can construct an irreducible metric with divisor D' such that $\beta_1, \beta_2, \beta_3 \notin \mathbf{Z}$: The metric $d\sigma_x^2 \in \text{Met}_1(S^2)$ obtained in Theorem 2.4 is the desired one.

$\mathcal{H}^3$ -reducible case:

First, we consider the case of $\mathcal{H}^3$ -reducible. In this case, β_1, β_2 and β_3 are integers and g in (2.1) is single-valued on S^2 , i.e., a rational function on $\mathbf{C} \cup \{\infty\}$.

Without loss of generality, we assume

$$(3.3) \quad \beta_1 \leq \beta_2 \leq \beta_3.$$

Let g be a rational function such that $d\sigma^2$ as in (2.1) with the divisor D in (2.6) satisfying (3.3). Then the ramification points of g are 0, 1 and ∞ whose orders are β_1, β_2 and β_3 respectively. By the Riemann-Hurwitz formula,

$$\deg g = \frac{1}{2}(\beta_1 + \beta_2 + \beta_3) + 1 \leq \frac{1}{2}\beta_1 + \beta_3 + 1 < (\beta_1 + 1) + (\beta_3 + 1)$$

holds. Then we have $g(p_1) \neq g(p_3)$, and similarly, $g(p_2) \neq g(p_3)$. Thus, by a suitable change as (2.2), we may assume $g(p_1) = g(0) \neq \infty$, $g(p_2) = g(1) \neq \infty$, and $g(p_3) = g(\infty) = \infty$. Under these assumptions, we can write

$$(3.4) \quad dg = c \frac{z^{\beta_1} (z-1)^{\beta_2}}{\prod_{j=1}^N (z-a_j)^2} dz, \quad \beta_3 = \beta_1 + \beta_2 - 2N,$$

where $c \neq 0$ is a constant, and $a_1, \dots, a_N \in \mathbf{C} \setminus \{0, 1\}$ are mutually distinct numbers. Conversely, if there exists g which satisfies (3.4), then we have $d\sigma^2 \in \text{Met}_1(S^2)$ with desired singularities. Computing residues at $z = a_1, \dots, a_N$, we have

Proposition 3.3. *Let β_1, β_2 and β_3 are positive integers satisfying (3.3). Then there exists $d\sigma^2 \in \text{Met}_1(S^2)$ with divisor D in (2.6) if and only if there exists a non-negative integer N and mutually distinct complex numbers $a_1, \dots, a_N \in \mathbf{C} \setminus \{0, 1\}$ which satisfy*

$$(3.5) \quad \beta_1 + \beta_2 - \beta_3 = 2N$$

and

$$(3.6) \quad \frac{\beta_1}{a_j} + \frac{\beta_2}{a_j - 1} - \sum_{k \neq j} \frac{2}{a_j - a_k} = 0, \quad (j = 1, \dots, N).$$

Remark. An $\mathcal{H}^3$ -reducible metric admits a three parameter space of deformation as in (2.5) which preserves the divisor D and the Schwarzian derivative. For a given triple $(\beta_1, \beta_2, \beta_3)$ as in Proposition 3.3, such a deformation space is determined uniquely. In fact, assume two metrics $d\sigma_1^2 = g_1^* d\sigma_0^2$ and $d\sigma_2^2 = g_2^* d\sigma_0^2$ have the same divisor D . Then both $\tilde{S}(d\sigma_1^2) = S(g_1)$ and $\tilde{S}(d\sigma_2^2) = S(g_2)$ are equal to (2.13). Then there exists $a \in \text{PSL}(2, \mathbf{C})$ such that $g_2 = a \star g_1$.

Observing the equation (3.6) for small N , one can easily see the following facts: If N in (3.5) is 0, trivially the metric exists.

When $N = 1$, (3.6) has the unique solution $a_1 = \beta_1/(\beta_1 + \beta_2) \neq 0, 1$.

Assume $N = 2$. Then $\beta_3 = \beta_1 + \beta_2 - 4$. By (3.3), this implies $4 \leq \beta_1 \leq \beta_2$. In this case, it is easy to show that the system of equations (3.6) has the unique solution up to the order of a_j 's.

Hence, we have

Corollary 3.4. *Let β_1, β_2 and β_3 are positive integers satisfying (3.3) and*

$$\beta_1 + \beta_2 - \beta_3 = 2N \quad (N = 0, 1 \text{ or } 2).$$

Then, there exists an $\mathcal{H}^3$ -reducible metric $d\sigma^2 \in \text{Met}_1(S^2)$ with the divisor D as in (2.6). Moreover, such a metric is unique up to three parameter deformation as in (2.5).

$\mathcal{H}^1$ -reducible case:

For $\mathcal{H}^1$ -reducible case, one of the β_j 's must be an integer because of Lemma 3.1. We assume that β_1 and β_3 are non-integral numbers, and β_2 is a positive integer:

$$\beta_1, \beta_3 \notin \mathbf{Z}, \quad \beta_2 \in \mathbf{Z}.$$

Then we can choose g as in (2.1) such that

$$(3.7) \quad g = z^\mu \varphi(z), \quad \mu \in \mathbf{R} \setminus \mathbf{Z},$$

where $\varphi(z)$ is a rational function. Here, such a normalization of g is unique up to the change $g \mapsto tg$ or $g \mapsto t/g$ for each non-zero constant t . Moreover,

$$(3.8) \quad (tg)^* d\sigma_0^2 \quad (t \in \mathbf{R}^+)$$

gives a non-trivial deformation of the metric preserving the divisor. This is the one parameter deformation as (2.5).

Let g be a function as in (3.7) such that $d\sigma^2$ as in (2.1) has the divisor D in (2.6). Replacing g with $1/g$, we may assume $\varphi(1) \neq \infty$ without loss of generality. Under this assumption, dg has zero of order β_2 at $z = 1$:

$$(3.9) \quad dg = cz^{\nu_1} \frac{(z-1)^{\beta_2}}{\prod_{j=1}^N (z-a_j)^2} dz \quad (\nu_1 = \mu - 1),$$

where $c \neq 0$ is a constant, and $a_1, \dots, a_N \in \mathbf{C} \setminus \{0, 1\}$ are mutually different numbers.

We denote the order of dg at $z = \infty$ by ν_3 :

$$(3.10) \quad \nu_3 = -\nu_1 - \beta_2 + 2N - 2.$$

So the following four cases occur.

- (a) $\nu_1 > -1$ and $\nu_3 > -1$. In this case, $\beta_1 = \nu_1$, $\beta_3 = \nu_3$. Hence $\beta_1 + \beta_2 + \beta_3 = 2N - 2$.
- (b) $\nu_1 < -1$ and $\nu_3 < -1$. In this case, $\beta_1 = -\nu_1 - 2$, $\beta_3 = -\nu_3 - 2$. Hence $\beta_1 - \beta_2 + \beta_3 = -2N - 2$.
- (c) $\nu_1 > -1$ and $\nu_3 < -1$. In this case, $\beta_1 = \nu_1$, $\beta_3 = -\nu_3 - 2$. Hence $\beta_1 + \beta_2 - \beta_3 = 2N$.
- (d) $\nu_1 < -1$ and $\nu_3 > -1$. In this case, $\beta_1 = -\nu_1 - 2$, $\beta_3 = \nu_3$. Hence $\beta_1 - \beta_2 - \beta_3 = -2N$.

For the cases (a) and (c), there exists a meromorphic function g on the universal cover of $\mathbf{C} \setminus \{0\}$ satisfying (3.9) if and only if

$$(3.11) \quad \frac{\beta_1}{a_j} + \frac{\beta_2}{a_j - 1} - \sum_{k \neq j} \frac{2}{a_j - a_k} = 0, \quad (j = 1, \dots, N)$$

holds, and for the cases (b) and (d), there exists g satisfying (3.9) if and only if

$$(3.12) \quad \frac{-\beta_1 - 2}{a_j} + \frac{\beta_2}{a_j - 1} - \sum_{k \neq j} \frac{2}{a_j - a_k} = 0, \quad (j = 1, \dots, N)$$

holds.

Then we have

Proposition 3.5. *Let β_1, β_3 be non-integral real numbers greater than -1 and β_2 a positive integer. Then there exists $d\sigma^2 \in \text{Met}_1(S^2)$ with divisor D in (2.6) if and only if one of the following occurs:*

- (1) *There exists non-negative integer N and mutually distinct complex numbers $a_1, \dots, a_N \in \mathbf{C} \setminus \{0, 1\}$ which satisfy $\beta_1 + \beta_2 + \beta_3 = 2N - 2$ and (3.11) holds.*
- (2) *There exists non-negative integer N and mutually distinct complex numbers $a_1, \dots, a_N \in \mathbf{C} \setminus \{0, 1\}$ which satisfy $\beta_1 - \beta_2 + \beta_3 = -2N - 2$ and (3.12) holds.*
- (3) *There exists non-negative integer N and mutually distinct complex numbers $a_1, \dots, a_N \in \mathbf{C} \setminus \{0, 1\}$ which satisfy $\beta_1 + \beta_2 - \beta_3 = 2N$ and (3.11) holds.*
- (4) *There exists non-negative integer N and mutually distinct complex numbers $a_1, \dots, a_N \in \mathbf{C} \setminus \{0, 1\}$ which satisfy $\beta_1 - \beta_2 - \beta_3 = -2N$ and (3.12) holds.*

Moreover, such a metric is unique up to the change in (3.8).

As well as $\mathcal{H}^3$ -reducible case, we classify the metrics for $N \leq 2$. It is easy to show the following lemma:

Lemma 3.6. *Let m and N be a positive integer, and ν a non-integral real number. Consider the following equations on $a_1, \dots, a_N$:*

$$(3.13) \quad \frac{\nu}{a_j} + \frac{m}{a_j - 1} - \sum_{\substack{k \neq j \\ 1 \leq k \leq N}} \frac{2}{a_j - a_k} = 0, \quad (j = 1, \dots, N).$$

Then

- (1) *If $N = 1$, the equation has the unique solution $a_1 = \nu/(\nu + m)$, which is different to 0 and 1.*
- (2) *If $N = 2$, the equation has a solution if and only if $m \neq 1$. The both solutions a_1, a_2 are distinct to 0 and 1 if $m \neq 1$.*

Using this Lemma, we have the following existence and non-existence result.

Corollary 3.7. *Let $\beta_2 = 1$, and β_1 and β_3 (> -1) be non-integral real numbers satisfying $\beta_1 + \beta_3 = 1$. Then there exists no metric $d\sigma^2 \in \text{Met}_1(S^2)$ with divisor D in (2.6).*

Proof. This is the case (1) in Proposition 3.5 for $N = 1$. Since $\beta_2 = 1$, the equation (3.11) has no solution because of Lemma 3.6. (q. e. d.)

Corollary 3.8. *Let β_1, β_3 be non-integral numbers (> -1) and β_2 an integer satisfying one of the following cases for some non-negative integer n :*

- (1) $\beta_1 + \beta_3 = 2n - 1, \beta_2 = 2n + 2N + 1$ ($N = 0, 1, 2$),
- (2) $\beta_1 + \beta_3 = 2n, \beta_2 = 2n + 2N + 2$ ($N = 0, 1, 2$),
- (3) $\beta_1 - \beta_3 = 2n + 1, \beta_2 = 2n + 2N + 1$ ($N = 0, 1, 2$),
- (4) $\beta_1 - \beta_3 = 2n, \beta_2 = 2n + 2N$ ($N = 0, 1, 2$).

Then there exists an $\mathcal{H}^1$ -reducible metric $d\sigma^2 \in \text{Met}_1(S^2)$ with divisor D in (2.6).

Proof. The first two cases follows from (2) of Proposition 3.5, and others from (4) of Proposition 3.5. (q. e. d.)

Appendix A.

In this appendix, we shall prove the following

Lemma A. *Let $a_j \in \text{SU}(2)$ ($j = 1, 2, 3$) be matrices satisfying $a_1 \cdot a_2 \cdot a_3 = \text{id}$. Then the following inequality holds:*

$$(A.1) \quad \cos^2 C_1 + \cos^2 C_2 + \cos^2 C_3 + 2 \cos C_1 \cos C_2 \cos C_3 \leq 1,$$

where $-e^{\pm iC_j}$ are the eigenvalues of the matrices a_j ($j = 1, 2, 3$). Moreover, equality holds if and only if a_1, a_2 and a_3 are diagonalizable at the same time.

Proof. We may assume that a_1 is diagonal without loss of generality. Then we may set

$$a_1 = \begin{pmatrix} -e^{iC_1} & 0 \\ 0 & -e^{-iC_1} \end{pmatrix}, \quad a_2 = \begin{pmatrix} p & -\bar{q} \\ q & \bar{p} \end{pmatrix} \quad (|p|^2 + |q|^2 = 1).$$

Since a_3^{-1} has the same eigenvalues as a_3 , we have

$$-2 \cos C_3 = \text{Tr}(a_1 \cdot a_2) = -pe^{iC_1} - \bar{p}e^{-iC_1}.$$

On the other hand, we have

$$\text{Trace } a_2 = p + \bar{p} = -2 \cos C_2.$$

Using the above two equations, we have

$$p = \pm i \frac{e^{\pm iC_1} \cos C_2 + \cos C_3}{\sin C_1}.$$

Thus we have

$$1 \geq |p|^2 = \frac{1}{\sin^2 C_1} (\cos^2 C_2 + \cos^2 C_3 + 2 \cos C_1 \cos C_2 \cos C_3).$$

This is equivalent to the desired inequality. Obviously, equality holds if and only if $|p| = 1$, that is, a_2 is diagonal. (q. e. d.)

Appendix B.

This is the appendix in [RUY1]. We attach it here for the sake of convenience.

Let Γ be a subgroup of $\text{PSU}(2)$.

In this appendix, we prove a property of a set of groups conjugate to Γ in $\text{PSL}(2, \mathbf{C})$ defined by

$$C_\Gamma := \{\sigma \in \text{PSL}(2, \mathbf{C}) \mid \sigma \cdot \Gamma \cdot \sigma^{-1} \subset \text{PSU}(2)\}.$$

The authors wish to thank Hiroyuki Tasaki for valuable comments on the first draft of the appendix.

If $\sigma \in C_\Gamma$, it is obvious that $a \cdot \sigma \in C_\Gamma$ for all $a \in \text{PSU}(2)$. So if we consider the quotient space

$$I_\Gamma := C_\Gamma / \text{PSU}(2),$$

the structure of the set C_Γ is completely determined. Define a map $\tilde{\phi} : C_\Gamma \rightarrow \mathcal{H}^3$ by

$$\tilde{\phi}(\sigma) := \sigma^* \cdot \sigma,$$

where $\mathcal{H}^3$ is the hyperbolic 3-space defined by $\mathcal{H}^3 := \{a \cdot a^* \mid a \in \text{PSL}(2, \mathbf{C})\}$. Then it induces an injective map $\phi : I_\Gamma \rightarrow \mathcal{H}^3$ such that $\phi \circ \pi = \tilde{\phi}$, where $\pi : C_\Gamma \rightarrow I_\Gamma$ is the canonical projection. So we can identify I_Γ with a subset $\phi(I_\Gamma) = \tilde{\phi}(C_\Gamma)$ of the hyperbolic 3-space $\mathcal{H}^3$. The following assertion holds.

Lemma B. *The subset $\phi(I_\Gamma)$ is a point, a geodesic line, or all $\mathcal{H}^3$.*

Proof. For each $\gamma \in \Gamma$, we set

$$C_\gamma := \{\sigma \in \text{PSL}(2, \mathbf{C}) \mid \sigma \cdot \gamma \cdot \sigma^{-1} \in \text{PSU}(2)\}.$$

Then we have

$$(B.1) \quad C_\Gamma := \bigcap_{\gamma \in \Gamma} C_\gamma.$$

The condition $\sigma \cdot \gamma \cdot \sigma^{-1} \in \text{PSU}(2)$ is rewritten as $\sigma^* \cdot \sigma \cdot \gamma = \gamma \cdot \sigma^* \cdot \sigma$. So we have

$$(B.2) \quad \tilde{\phi}(C_\gamma) = \mathcal{H}^3 \cap Z_\gamma,$$

where Z_γ is the center of $\gamma \in \Gamma$.

Assume $\gamma \neq \pm \text{id}$. If γ is a diagonal matrix, it can easily be checked that Z_γ consists of diagonal matrices in $\text{PSL}(2, \mathbf{C})$. Since any $\gamma \in \Gamma$ can be diagonalized by a matrix in $\text{SU}(2)$, we have $Z_\gamma = \{\exp(zT) \mid z \in \mathbf{C}\}$, where $T \in \mathfrak{su}(2)$ is so chosen that $\gamma = \exp(T)$. Hence we have

$$(B.3) \quad \tilde{\phi}(C_\gamma) = \mathcal{H}^3 \cap Z_\gamma = \exp(i\mathbf{R}T),$$

because $\exp(i\mathfrak{su}(2)) = \mathcal{H}^3$.

Suppose now that Γ is not abelian. Then there exist $\gamma, \gamma' \in \Gamma$ such that $\gamma \cdot \gamma' \neq \gamma' \cdot \gamma$. Set $\gamma = \exp(T)$ and $\gamma' = \exp(T')$, where $T, T' \in \mathfrak{su}(2)$. Then we

have $i\mathbf{R}T \cap i\mathbf{R}T' = \{0\}$. It is well-known that the restriction of the exponential map $\exp|_{i\mathfrak{su}(2)}: i\mathfrak{su}(2) \rightarrow \mathcal{H}^3$ is bijective. Hence we have

$$\tilde{\phi}(C_\gamma) \cap \tilde{\phi}(C_{\gamma'}) = \exp(i\mathbf{R}T) \cap \exp(i\mathbf{R}T') = \{\text{id}\}.$$

By (B.1), (B.2) and (B.3), we have

$$\phi(I_\Gamma) = \{\text{id}\} \quad (\text{if } \Gamma \text{ is not abelian}).$$

Next we consider the case Γ is abelian. If $\Gamma \subset \{\pm \text{id}\}$, then obviously

$$\phi(I_\Gamma) = \mathcal{H}^3.$$

Suppose $\Gamma \not\subset \{\pm \text{id}\}$. Then there exists $\gamma \in \Gamma$ such that $\gamma \neq \pm \text{id}$. We set $\gamma = \exp T$ ($T \in \mathfrak{su}(2)$). Since $\exp(\mathbf{R}T)$ is a maximal abelian subgroup containing γ , we have $\Gamma \subset \exp(\mathbf{R}T)$. Then by (B.3), we have

$$\phi(I_\Gamma) = \exp(i\mathbf{R}T).$$

(q. e. d.)

References.

- [B] R. Bryant, *Surfaces of mean curvature one in hyperbolic space*, Astérisque **154–155** (1987), 321–347.
- [CL] W. Chen and C. Li, *What kinds of singular surfaces can admit constant curvature?*, Duke Math. J., **78** (1995) 437–451.
- [LT] F. Luo and G. Tian, *Liouville equation and spherical convex polytopes*, Proc. Amer. Math. Soc., **116** (1992), 1119–1129.
- [H] Y. Hattori, in preparation.
- [RUY1] W. Rossman, M. Umehara and K. Yamada, *Irreducible constant mean curvature 1 surfaces in hyperbolic space with positive genus*, to appear in Tôhoku Math. J.
- [RUY2] W. Rossman, M. Umehara and K. Yamada, *Metrics with conical singularities and CMC-1 surfaces of low total curvature in hyperbolic space*, in preparation.
- [Sm] A. J. Small, *Surfaces of Constant Mean Curvature 1 in H^3 and Algebraic Curves on a Quadric*, Proc. Amer. Math. Soc. **122** (1994), 1211–1220.
- [T1] M. Troyanov, *Metric of constant curvature on a sphere with two conical singularities*, in “Differential Geometry”, Lect. Notes in Math. vol. 1410, Springer-Verlag, (1989), 296–306.
- [T2] M. Troyanov, *Prescribing curvature on compact surfaces with conical singularities*, Trans. Amer. Math. Soc. **324** (1991), 793–821.
- [UY1] M. Umehara and K. Yamada, *Complete surfaces of constant mean curvature-1 in the hyperbolic 3-space*, Ann. of Math. **137** (1993), 611–638.
- [UY2] M. Umehara and K. Yamada, *Surfaces of constant mean curvature- c in $H^3(-c^2)$ with prescribed hyperbolic Gauss map*, Math. Ann. **304** (1996), 203–224.
- [UY3] M. Umehara and K. Yamada, *A duality on CMC-1 surface in the hyperbolic 3-space and a hyperbolic analogue of the Osserman Inequality*, Tsukuba J. Math. **21** (1997), 229–237.

MASAAKI UMEHARA: Department of Mathematics, Graduate School of Science, Osaka University, Toyonaka, Osaka 560, Japan. *E-mail*: umehara@math.wani.osaka-u.ac.jp

KOTARO YAMADA: Department of Mathematics, Faculty of Science, Kumamoto University, Kumamoto 860, Japan. *E-mail*: kotaro@gpo.kumamoto-u.ac.jp